

Entomofauna

ZEITSCHRIFT FÜR ENTOMOLOGIE

Band 25, Heft 8: 117-132

ISSN 0250-4413

Ansfelden, 10. August 2004

**Contribution to the knowledge of Italian cleptoparasitic Bees. X.
The genus *Sphecodes* LATREILLE, '*pinguiculus* PÉREZ' group,
with description of new species
(Hymenoptera, Apoidea, Halictidae)**

Vittorio NOBILE & Giuseppe Fabrizio TURRISI

Abstract

The authors propose the institution of a "*pinguiculus* PÉREZ" group, belonging to the genus *Sphecodes* LATREILLE, and describe 7 new species of this group: 3 from peninsular Italy (*S. campadellii* sp. nov., *S. combai* sp. nov., *S. banaszaki* sp. nov.) and 4 from Sicily (*S. marcellinoi* sp. nov., *S. walteri* sp. nov., *S. iosephi* sp. nov., *S. tomarchioi* sp. nov.). A key for the identification of the ♂♂ of the W-palaeartic species of the genus *Sphecodes pinguiculus* group is provided.

Key words: Italy, *Sphecodes pinguiculus* group, Halictidae, new species.

Zusammenfassung

Die Verfasser schlagen die Errichtung einer "*pinguiculus* PÉREZ-Gruppe" in der Gattung *Sphecodes* vor mit der Präsentation von 7 neuen Arten: 3 von der Halbinsel Italien (*S. campadellii* sp. nov., *S. combai* sp. nov., *S. banaszaki* sp. nov.) und 4 von Sizilien (*S. marcellinoi* sp. nov., *S. walteri* sp. nov., *S. iosephi* sp. nov., *S. tomarchioi* sp. nov.). Es wird ein Bestimmungsschlüssel für die ♂♂ der westpaläarktischen Arten der *Sphecodes pinguiculus*-Gruppe angeboten.

Introduction

In the genus *Sphecodes* LATREILLE, 1804, the "*pinguiculus* PÉREZ" group, for which the institution is proposed, is characterised by the fact that the small white setae which

cover the posterior part of the male antennae of the species are regularly and uniformly distributed on the articles of the flagellum, rather than densely packed together in particular areas at the base of these articles, as it is the case in the majority of the other species of Sphecodini. The dislocation and the size of these setaceous areas, in fact, are specific and constitute a good aid for the distinction of the various species. Another common character of the males of this group is that the parameral plates of the genital capsule, dorsally and longitudinally, have a deep, wide groove (unlike other groups in which the parameral plates appear full and compact). This character is also shared by other groups, but the fact that the *pinguiculus* group comprises species with specimens of small dimension, never exceeding 7 mm, is noteworthy. A further characteristic of the group (with the exception of *Sphecodes campadellii* sp. n.) is that the tarsi, the proximal and distal parts of the tibiae and the distal part of femora, are yellow-reddish: lighter in colour than the rest of the legs; hairs on the surface of the body white as generally.

In the Western sector of the Palearctic region, the genus *Sphecodes*, "*pinguiculus*" group, has been known for only two species: *S. pinguiculus* PÉREZ, 1903 and *S. intermedius* BLÜTHGEN, 1923, both present also in Europe; of these, however, only the former was part of the Italian fauna. In fact, WARNCKE (1992) reported it in Sardinia with the subspecies *S. pinguiculus sareptensis* MEYER, 1922 and NOBILE & CAMPADELLI (1998) reported it also for Sicily with the same subspecies.

Material and methods

The recent opportunity to examine some collections both public and private of Apoidea Sphecodini, allows us to increase greatly the number of species of the *pinguiculus* group not only in Sicily, but also to report the presence of some of them (the first time for this group) also in peninsular Italy. In fact, the species found are all new for science, 3 species from Italy and 4 from Sicily.

The studied material was generously provided by public institutions and private collections, for which we are grateful:

- Guido CAMPADELLI, Institute of Entomology "G. Grandi", University of Bologna.
- Mario COMBA (Cecchina, Rome).
- Salvatore TOMARCHIO (Catania).

Results and discussion

Sphecodes pinguiculus PÉREZ, 1903 ssp. *sareptensis* MEYER, 1922 (fig. 8)

Sphecodes sareptensis MEYER, 1922.- Arch. Naturg. 88A (8): 170-171 (Russia).

Sphecodes pinguiculus ssp. *sareptensis*: WARNCKE 1992: 32, 52, 61 (Sardinia: Cagliari).

Sphecodes pinguiculus ssp. *sareptensis*: NOBILE & CAMPADELLI 1998: 96 (Sicily).

Mediterranean-turanian-central Asian distribution, extending to the Canary Islands but excluding Spain where the nominal subspecies is present.

Sphecodes campadellii sp. nov. (fig. 1)

Holotypus ♂. Italy: Emilia Romagna, Zocca (Modena), 8.VIII.1962 (coll. Institute of Entomology "G. Grandi", University of Bologna).

♂. Length 5.0 mm. Head black, antennal scape dark brown; antennal flagellum brown

posteriorly, dark brown anteriorly; mandibles black. Thorax black, tegulae brownish. Wings slightly but uniformly dark, with blackish stigma and brown veins; legs brown, tibial spines whitish. Abdomen brown although not uniformly so, with a darker tone on disc of tergites.

Head roundish, a little wider than long; antennae short, robust and slightly knotted: the median antennomeres (4th-11th) have a wide impression on posterior face, not very deep, which can occupy up to 3/4 of the surface of the antennomeres, giving the flagellum a knotted appearance; 2nd antennomere 0.5 times shorter than wide, 3rd antennomere 1.2 times longer than wide, 4th antennomere as long as the 3rd, the remaining antennomeres a little longer than wide; clypeus almost flat; punctuation of the clypeus large, irregular, deep and dense (distance between the punctures generally equal to the diameter of one puncture); frons shagreened with coarse punctures, irregular, dense and deep (distance between the punctures less than the diameter of one puncture); vertex with small and superficial punctures, irregularly placed and sparse (distance between punctures 1-3 times the diameter of one puncture).

Mesothorax smooth, with large punctuation, deep, irregular and sparse on disc, denser on margins (distance between the punctures generally equal to the diameter of one puncture); scutellum with large punctures, deep, irregularly spaced and sparser than those of the thorax disc (distance between punctures greater than the diameter of one puncture); postscutellum shagreened with an irregular structure, with medium-large punctures, very dense; median field of the propodeum as long as the scutellum, almost bluntly triangular in shape, with the distal corner replaced by a cavity; margins of the median field distinct, forming a ridge, raised, and its internal structure is irregular honey-comb like; lateral and posterior parts of the propodeum with large unequal alveoli with a honey-comb structure, similar to that of the median field; mesopleurae shagreened, with an irregular honey-comb structure. Wings with the first discoidal vein terminating just before the second cubital vein; second cubital cell longer than wide.

Abdomen very shiny; T₁ with few small punctures, superficial and sparse, irregularly spaced; T₂₋₄ with less sparse punctures; successive tergites with not very evident punctures, substituted by wrinkles. Distal part of all the tergites without punctures, transparent and shiny.

Face with medium length setae, sparse (the underlying punctures are visible); scape with medium length setae and averagely dense; vertex with medium length sparse setae; postgena with medium sparse setae; mesothorax and scutellum with short, sparse setae; postscutellum with long sparse setae, propodeum with medium, sparse setae; mesopleurae with medium sparse setae, ventral parts of the thorax with short, sparse setae; abdomen, dorsally, with very sparse short setae, tergites with short, sparse lashes, sternites with short, sparse setae together with short, sparse lashes.

♀-unknown.

Derivatio nominis: We dedicate this species to the late colleague and friend Guido CAMPADELLI (University of Bologna).

Sphecodes combai sp. nov. (fig. 2)

Holotypus ♂. Italy: Latium, Albani hills, Cecchina, Poggio Ameno, 280 m, I.X.1990. M. COMBA leg., on *Hedera helix* L. (Zoologische Staatssammlung München).

♂. Length 4.0 mm. Head black, antennal scape dark brown; flagellum posteriorly brown, slightly darker anteriorly; mandibles brown at base and light-brown for the rest. Thorax black, tegulae brownish. Wings slightly darkened, with stigma and veins brown; legs brown except the tarsi and the proximal and distal tips of the tibiae, which are yellow-orange; tibial spines yellow-orange. Abdomen brown, not uniformly.

Head roundish, a little wider than long; antennae short and robust; 2nd antennomere 0.6 times shorter than wide, 3rd antennomere 1.2 times longer than wide, 4th antennomere as long as the 3rd, the remaining antennomeres a little longer than wide; clypeus slightly curved. Punctuation on clypeus large, irregular, deep and dense (distance between the punctures generally equal to the diameter of one puncture); frons with largish punctuation, almost regular, dense and deep (distance between the punctures less than the diameter of one puncture); vertex with largish punctures, superficial and sparse, irregularly placed (distance between punctures 1-3 times the diameter of one puncture).

Mesothorax smooth, with large punctuation, deep, irregular and sparse on the disc, denser at margins (distance between the punctures generally equal to the diameter of one puncture); scutellum with large punctures, deep, moderately dense, irregularly placed (distance between punctures generally less than the diameter of one puncture); postscutellum shagreened with an irregular structure, with medium large punctures, very dense; median field of propodeum as long as the scutellum, half-moon shaped, with distinct margins, forming a very raised ridge: inside, the alveoli are delimited by a subparallel ridge, with a radial course and by few transversal ridges. Lateral and posterior parts of the propodeum with large alveoli with an irregular honey-comb structure; mesopleurae shagreened, with an irregular honey-comb structure, with a smooth ridge. Wings with the first discoidal, almost interstitial vein, terminating just before the second cubital vein; second cubital cell longer than wide.

Abdomen shiny; disc of T₁ with few small sparse punctures (distance between the punctures 1-3 times the diameter of one puncture); the declivous part of T₁ with punctures even more sparse; T₂₋₃ disc with denser and more regular punctuation (distance between punctures about a diameter of one puncture); the remaining tergites with not very evident punctures, replaced by wrinkles; distal part of all the tergites without punctures, transparent and shiny.

Face with medium length setae, sparse (the underlying punctures are visible); scape with medium setae and not very dense; vertex with medium length sparse setae; postgena with medium sparse setae; mesothorax and scutellum with medium, sparse setae; postscutellum with long setae averagely dense; propodeum with medium, sparse setae; mesopleurae with medium sparse setae; ventral parts of the thorax with medium, not very dense setae; abdomen, dorsally, with very sparse short setae; tergites with short, sparse setae; T₃₋₇ with short lashes; sternites with short, sparse setae together with short, sparse lashes.

♀ unknown.

Derivatio nominis: We dedicate this species to Mario COMBA from Cecchina (Rome).

Sphecodes banaszaki sp. nov. (fig. 3)

Holotypus ♂. Italy, Lazio, Albani hills, Communal forest of Albano, 480 m, 23.VI. 1995, M. COMBA leg. (Zoologische Staatssammlung München).

♂. Length 5.0 mm. Head and antennal scape black, 1st and 2nd antennomeres dark

brown; remaining part of the flagellum light brown posteriorly, dark brown anteriorly; mandibles black at base, yellow medially and red apically. Thorax black, tegulae brownish; wings hyaline, with stigma black and veins blackish; legs black, except the tarsi, the proximal and distal parts of the tibiae and the distal parts of the femura, which are orange; tibial spines yellow-orange. Abdomen dorsally black, with reddish transparencies on the distal margins of T_{1-2} and on the proximal margins of T_{2-3} .

Head roundish, a little wider than long; antenna short and robust, 2nd antennomere 0.6 times shorter than wide, 3rd antennomere 1.2 times longer than wide, 4th antennomere 1.1 times longer than wide, the remaining antennomeres slightly longer than wide; clypeus curved, with large punctuation, irregular, deep, dense (distance between the punctures generally equal to the diameter of one puncture); frons shagreened, with large punctuation, irregular, dense and deep (distance between the punctures less than the diameter of one puncture); vertex shagreened, with large punctures, deep, sparse and irregularly placed (distance between the punctures 1-3 times the diameter of one puncture).

Mesothorax and scutellum shagreened, with large punctuation, deep, irregular and not very dense (distance between the punctures generally less than the diameter of one puncture); postscutellum shagreened, with large punctures, very dense; median field of the propodeum as long as the scutellum, half-moon shaped, with clear margins, ridge very raised, with a very irregular honey-comb structure inside; lateral and posterior parts of the propodeum with irregular honey-comb structure, with alveoli smaller than those of the median field; mesopleurae shagreened, with irregular honey-comb structure, with raised ridges. Wings with the first discoidal vein terminating at 1/3 before the second cubital vein; second cubital cell a little longer than wide.

Abdomen opaque; surfaces of all the tergites shagreened; disc of T_{1-2} with medium small punctures and not very dense (distance between the punctures 1 - 1.5 times the diameter of one puncture); disc of T_3 with punctuation a little more sparse and more superficial than the preceding tergites; distal part of all the tergites without punctures, transparent, not very shiny.

Face with setae of medium length, averagely dense (the underlying punctures are visible); antennal scape with medium setae and not very dense; vertex with sparse, medium length setae; postgena with medium setae, not very dense; mesothorax and scutellum with short and sparse setae; postscutellum with long setae not very dense; propodeum with medium, sparse setae; mesopleurae with medium, not very dense setae; ventral parts of the thorax with medium length setae not very dense; abdomen, dorsally, with very sparse short setae and with short lashes on T_{3-7} , not very sparse; all the sternites with short setae and short, sparse lashes.

♀ unknown.

Derivatio nominis: We dedicate this species to Józef BANASZAK from the University of Bydgoszcz (Poland).

Sphecodes marcellinoi sp. nov. (fig. 4)

Holotypus ♂. Italy, Sicily: Mount Etna, Piano Vetore, 1700 m (Ragalna), 2.V.1999, S. TOMARCHIO leg. (Zoologische Staatssammlung München).

♂. Length 4.2 mm. Head, antennal scape and 1st and 2nd antennomeres black; remaining flagellum light brown posteriorly, slightly darker anteriorly; mandibles black at base, me-

dially yellow and red apically. Thorax black; tegulae hyaline, transparent; wings hyaline with dark stigma and brown veins; legs black, except tarsi, the proximal and distal tips of tibiae, the distal tips of femura and the tibial spines, yellow-orange in colour. Abdomen dorsally black, except the distal margins of T₁₋₃ and the proximal margins of T₂₋₄, reddish and transparent.

Head roundish, a little wider than long; 2nd antennomere 0.5 times shorter than wide; 3rd antennomere 1.3 times longer than wide; 4th antennomere as long as 3rd, the remaining antennomeres slightly longer than wide (almost square); clypeus slightly curved; large punctuation of clypeus, irregular, deep and dense (distance between the punctures generally equal to the diameter of one puncture); frons with fine punctures, almost regular, sparse and superficial (distance between the punctures about two punctures diameters); vertex with small punctures, superficial and sparse, irregularly placed (distance between the punctures about two punctures diameters).

Mesothorax with large punctuation, deep, irregular and not very dense (distance between the punctures generally less than the diameter of one puncture); scutellum with large punctures, deep and sparse, irregularly placed (distance between the punctures generally about one diameter); postscutellum shagreened, with irregular honey-comb structure; median field of the propodeum a little shorter than the scutellum, half-moon shaped with clear margins (with a very raised ridge), with grooves and subparallel longitudinal ridges; lateral and posterior parts of the propodeum with large, unequal alveoli and a honey-comb structure; mesopleurae shagreened with irregular honey-comb structure. Wings with the first discoidal vein terminating at 1/4 before the second cubital vein; second cubital cell longer than wide.

Abdomen averagely shiny; disc, and declivous part of T₁, with medium small sparse punctures, irregularly placed (distance between punctures 1-2 diameters of one puncture); disc of T₂ with similar punctures as T₁ but slightly denser; discs of the remaining tergites with punctuation and shagreening; distal part of all the tergites without punctures, transparent and not very shiny.

Face with medium length setae, dense (hiding the underlying punctuation); scape with medium, sparse setae; vertex with long, sparse setae; postgena with medium length not very dense setae; mesothorax with medium, sparse setae; scutellum with short, sparse setae; postscutellum with long averagely dense setae; propodeum with short, sparse setae; mesopleurae with medium length not very dense setae; ventral parts of the thorax with dense medium length setae; abdomen, dorsally, with very sparse short setae, T₄₋₇ with lashes; sternites with sparse, short setae; short lashes on all the sternites.

♀ unknown.

Derivatio nominis: We dedicate this species to Italo MARCELLINO, Arachnologist, of the University of Catania.

Sphcodes walteri sp. nov. (fig. 5)

Holotypus ♂. Italy, Sicily, Mount Etna south slope, c.da Milia, 1500 m (Ragalna), 10.VIII.1999, S. TOMARCHIO leg. (Zoologische Staatssammlung München).

♂. Length 5.0 mm. Head, antennal scape and 1st and 2nd antennomeres black; remaining flagellum dark brown posteriorly, much darker, almost black, anteriorly; mandibles black at base, medially yellow and red apically. Thorax black, tegulae hyaline; wings hyaline

with stigma and veins brownish; legs black, except tarsi, the proximal and distal tips of tibiae and the distal tips of femura, brown in colour; tibial spines whitish. Abdomen, dorsally, black, except the distal margin of $T_{1,3}$ and the proximal margin of $T_{2,3}$, reddish and transparent.

Head wider than long; 2nd antennomere 0.6 times shorter than wide, 3rd antennomere 1.3 times longer than wide, 4th antennomere as long as 3rd, the remaining antennomeres a little longer than wide; clypeus notably arcuate; punctuation of the clypeus large, irregular, deep and dense (distance between the punctures generally equal to the diameter of one puncture); frons with irregular punctuation, irregular, deep and not very dense (distance between the punctures equal to 1-2 diameters of one puncture); vertex with irregular punctures, not very deep and not dense, irregularly placed (distance between the punctures equal to 1-3 diameters of one puncture).

Mesothorax smooth with large punctuation, deep, irregular and dense on disc, denser on margins (distance between the punctures generally less than a puncture diameter); scutellum with large punctures, deep and dense, irregularly placed as on disc of mesothorax; postscutellum shagreened, with large, deep punctures, very dense and irregularly placed; median field of propodeum as long as scutellum, half-moon shaped, with clear margins, forming a prominent ridge, and with an internal irregular honey-comb structure; lateral and posterior parts of the propodeum with large alveoli with an irregular honey-comb structure, similar to that of the median field; mesopleurae shagreened with an irregular honey-comb structure. Wings with the first discoidal vein terminating just in front of the second cubital vein; second cubital cell longer than wide.

Abdomen shiny; T_1 with not very dense punctures, irregularly placed (distance between the punctures equal to 1-2 diameters of one puncture); $T_{2,3}$ with deeper punctures, a little denser than on T_1 ; the remaining tergites with not very evident punctuation, substituted by roughness; distal part of all the tergites without punctures, transparent and shiny.

Face with medium length setae, not very dense (they do not hide the underlying punctures); antennal scape with medium length setae, average density; vertex with long, sparse setae; postgena with short medium length setae, not very dense; mesothorax and scutellum with short setae, generally dense; postscutellum with long, not very dense setae; propodeum with short, sparse setae; mesopleurae with medium setae, generally dense; ventral parts of thorax with medium length setae, generally dense; abdomen, dorsally, with very sparse, short setae, and sparse lashes on all the tergites, including the declivous part of T_1 ; all the sternites with setae and sparse, short lashes.

♀ unknown.

Derivatio nominis: We dedicate this species to Walter DE LEONARDIS, Palinologist at the University of Catania.

Sphcodes iosephi sp. nov. (fig. 6)

Holotypus ♂. Italy, Sicily, Mount Etna, south slope, c.da Milia, 1400 m (Ragalna), 10.VIII.1999, S. TOMARCHIO leg. (Zoologische Staatssammlung München).

♂. Length 5.0 mm. Head, antennal scape and successive two antennomeres black; remaining part of flagellum dark brown posteriorly, notably darker anteriorly; mandibles black. Thorax black, tegulae brownish. Wings slightly and uniformly darkened, with stigma and veins brownish. Legs black, except tarsi, the proximal and distal tips of tibiae

and the distal tips of femura, brownish in colour; tibial spines whitish. Abdomen dorsally black, but with the distal margin of T₁₋₃ and the proximal margin of T₂₋₄ brownish and transparent.

Head roundish, a little wider than long; antenna robust; the antennomeres, at about 2/3 from base, arc bulging like a small barrel, giving the flagellum a knotty appearance; 2nd antennomere 0.6 times shorter than wide, 3rd antennomere 1.4 times longer than wide, 4th antennomere as long as 3rd, the remaining antennomeres longer than wide; clypeus arcuate; punctuation of the clypeus large, irregular, dense and deep (distance between the punctures generally equal to the diameter of one puncture); frons shagreened with largish punctures, irregular, very dense and deep (distance between the punctures notably less than the diameter of one puncture); vertex shagreened; on it the large, dense and deep punctuation coexists with the transverse prominent ridge structure, which is similar to the structure of the median field.

Mesothorax slightly shagreened, with large, deep, irregular, dense punctuation on disc, slightly denser on margins (distance between punctures generally less than the diameter of one puncture); scutellum with deep punctures, irregularly placed, less dense and smaller than those of the mesothorax disc (distance between punctures about as the diameter of one puncture); postscutellum shagreened with an irregular structure, with large, deep and very dense punctures; median field of the propodeum slightly longer than scutellum, half-moon shaped, with clear margins, with well raised ridges, and inside having an irregular honey-comb structure; lateral and posterior parts of the propodeum with large alveoli with an irregular honey-comb structure similar to that of the median field; mesopleurae slightly shagreened with an irregular honey-comb structure. Wings with the first discoidal vein terminating at 1/5 before the second cubital vein; second cubital cell almost as wide as long.

Abdomen shagreened, opaque; disc of T₁ with small, superficial punctures, irregularly placed and sparse (distance between punctures 1-3 diameters of one puncture); disc of T₂₋₃ with punctuation slightly denser and deeper (distance between punctures 1-2 diameters of one puncture); remaining tergites with punctuation not very clear, substituted by roughness; distal part of all the tergites without punctures and transparent.

Face with medium length setae, and averagely dense (the underlying punctuation is visible); antennal scape with medium length setae and rather dense; vertex with medium length, sparse setae; postgena with short, medium length not very dense setae; mesothorax and scutellum with short and sparse setae; postscutellum with long, not very dense setae; propodeum with short, sparse setae; mesopleurae with medium length setae which are generally dense; the ventral parts of the thorax with medium length setae and averagely dense; abdomen dorsally with short, sparse setae and with lashes on all the tergites, though longer and denser on the last four; sternites with short, dense setae.

♀ unknown.

Derivatio nominis: We dedicate this species to Giuseppe NOBILE, son of the first author.

Sphecodes tomarchioi sp. nov. (fig. 7)

Holotypus ♂. Italy, Sicily, Mount Etna, south slope, c.da Milia, 1400 m (Ragalna), 10.VIII.1999, S. TOMARCHIO leg. (Zoologische Staatssammlung München).

♂. Length 4.6 mm. Head, antennal scape and 1st and 2nd antennomeres black; remaining part of the flagellum dark brown posteriorly, black anteriorly; mandible black at base, medially yellow, red apically. Thorax black, tegulae hyaline. Wings slightly and uniformly darkened, with stigma and veins brownish; legs dark brown except the tarsi, the proximal and distal tips of tibiae and the distal tips of femura light brown in colour; tibial spines whitish. Abdomen dorsally black, except the distal hollow of the tergites, brownish and transparent.

Head wider than long; antenna short and robust; 2nd antennomere 0.5 times shorter than wide, 3rd antennomere 1.4 times longer than wide, 4th antennomere 1.3 times longer than wide, remaining antennomeres longer than wide; clypeus arcuate; punctuation of the clypeus large, irregular, deep and very dense (distance between punctures less than the diameter of one puncture); frons with largish punctures, irregular, very dense and deep (distance between the punctures notably less than the diameter of one puncture); vertex shagreened with largish punctures, dense and deep, irregularly placed (distance between the punctures less than the diameter of one puncture).


Mesothorax smooth, with large punctuation, deep, irregular and dense (distance between the punctures generally less than the diameter of one puncture); scutellum shagreened, with large punctures on the disc, less in the periphery, deep, irregularly placed and a little denser than on disc of mesothorax (distance between the punctures generally less than the diameter of one puncture); postscutellum shagreened with an irregular structure, with medium large punctures, deep and very dense; median field of the propodeum a little longer than the scutellum, half-moon shaped, with clear margins, forming a very raised ridge with an internal irregular honey-comb structure; lateral and posterior parts of the propodeum with an irregular honey-comb structure and alveoli as large as those of the median field; mesopleurae shagreened with an irregular honey-comb structure. Wings with the first discoidal vein terminating at 1/4 before the second cubital vein; second cubital cell a little longer than wide.

Abdomen shiny; T₁ with punctures small, superficial, irregularly distributed and sparse (distance between the punctures 2-4 diameters of one puncture); T₂ with punctures less superficial and less sparse than on T₁ (distance between the punctures equal to 2-3 times the diameter of one puncture); T_{3,4} with punctures slightly denser than T₂, but less deep and less evident; the remaining tergites rough; distal part of the tergites without punctures, transparent and shiny.


Face with medium length setae, generally dense (the underlying punctures are still visible); scape with medium length setae and generally dense; vertex with medium length, sparse setae; postgena with medium length setae and not very dense; mesothorax and scutellum with short, sparse setae; postscutellum with long, not very dense setae; propodeum with short, sparse setae; mesopleurae with medium length setae and generally dense; ventral parts of thorax with medium length setae and generally dense; abdomen, dorsally with short, sparse setae and short, sparse lashes; sternites with short, sparse setae.

♀ unknown.




Derivatio nominis: We dedicate this species to Salvatore TOMARCHIO, friend and colleague of the authors.




1 - *Sphec. campadellii* sp. n.





2 - *Sphec. combai* sp. n.





3 - *Sphec. banaszaki* sp. n.





4 - *Sphec. marcellinoi* sp. n.








5 - *Sphec. walteri* sp. n.



6 - *Sphec. iosephi* sp. n.



7 - *Sphec. tomarchioi* sp. n.



8 - *Sphec. pingiculus* Pérez

Drawings by M.C. LA ROCCA, except for fig. 8, drawn from WARNCKE (1992).

Remarks

The small Sphecodini, as those of the genus *Sphecodes pinguiculus* group, are clepto-parasitic of collecting bees of the tribes Colletini, Andrenini and Halictini (RADCHENKO & PESENKO 1994), and also small in size. As there are about 200 species of collecting bees present in Italy with these characteristics, it is believed that probably, in the future, other species of the *pinguiculus* group and other species of other groups of Sphecodini will be found.

Key for the identification of the ♂♂ of the W-palaeartic species of the *Sphecodes* LATREILLE, "*pinguiculus* PÉREZ" group

- | | | |
|---|--|--|
| 1 | Third article of the flagellum twice as long as wide; length of the body 6-7 mm | <i>S. intermedius</i> BL. |
| | - Third article of the flagellum shorter; length of the body shorter | 2 |
| 2 | Mandibles black | 3 |
| | - Mandibles with three colours: black at base, red distally and orange medially | 4 |
| 3 | Surface of T ₁ very shiny, smooth; punctuation very sparse; length of the body 5 mm (genital capsule fig. 1) | <i>S. campadellii</i> sp. nov. |
| | - Surface of T ₁ opaque due to the shagreening, punctuation dense; length of the body 5 mm (genital capsule fig. 6) | <i>S. iosephi</i> sp. nov. |
| 4 | Surface of T ₁ opaque due to the shagreening; length of the body 5 mm (genital capsule fig. 3) | <i>S. banaszaki</i> sp. nov. |
| | - Surface of T ₁ shiny and smooth | 5 |
| 5 | T ₁₋₃ black and opaque apically, dark red and transparent distally; length of the body 4,6 mm (genital capsule fig. 7) | <i>S. tomarchioi</i> sp. nov. |
| | - Tergites of different colour | 6 |
| 6 | Disc of the tergites brown, with distal margins transparent, light brown in colour; length of the body 4 mm (genital capsule fig. 2) | <i>S. combai</i> sp. nov. |
| | - Abdomen of different colour; length of the body also greater | 7 |
| 7 | Antenna long, third antennomere 1.5 times longer than wide; body appearance robust; length of the body 4 - 5.5 mm (genital capsule fig. 8) | <i>S. pinguiculus</i> PÉREZ |
| | - Antenna shorter; third antennomere shorter; appearance of the body graceful | 8 |
| 8 | Posterior part of the flagellum and tarsi orange in colour; length of the body 4.2 mm (genital capsule fig. 4) | <i>S. marcellinoi</i> sp. nov. |
| | - Posterior part of the flagellum and tarsi brown in colour; length of the body 5 mm (genital capsule fig. 5) | <i>S. walteri</i> sp. nov. |

Bibliography

- NOBILE, V. & CAMPADELLI, G. - 1998. Il genere *Sphecodes* LATREILLE, 1804 in Italia (Hymenoptera, Apoidea, Halictidae). - Boll. Ist. Ent. "G. Grandi", Univ. Bologna 52: 85-103.
- RADCHENKO V.G. & PESENKO, Y.A. - 1994. Biology of Bees (Hymenoptera, Apoidea). - Russian Acad. Sciences, Zool. Institute, St. Petersburg: 350 pp. (in Russian).
- WARNCKE, K. - 1992. Die westpalacarktischen Arten der Bienengattung *Sphecodes* LATR. (Hymenoptera, Apidae, Halictinae). - Ber. Naturf. Ges., Augsburg 52: 9-64.

Authors' address:

Dr. Vittorio NOBILE / Dr. Giuseppe Fabrizio TURRISI
University of Catania
Department of Animal Biology "Marcello La Greca"
via Androne 81
I-95124 Catania, Italy
e-mail Dr. N. VITTORIO: nobilevittorio@tin.it
e-mail Dr. G.F. TURRISI: turrisifabrizio@yahoo.it